

6.0 DISCUSSION

6.1 Study Results

Seasonal and annual variability are essential factors in determining trends and are the reason it is important for volunteers to provide information on atmospheric weather conditions and qualitative data. Some of the lakes in the program have more than twenty years of historical data while others only have a handful of years. Weather plays an important role in the variability of data collected. As mentioned before, 2011 - 2013 were colder and wetter than average. In 2014 - 2016, many program lakes were much warmer than they were historically. In 2017 and 2018 we had hot, dry summers with a lot of forest fires that could affect lake evaporative rates, temperature, clarity, and nutrient levels. It will be important to continue to maintain consistency in data collection in the future so that trends become apparent in lieu of annual variation.

Nutrient concentrations are generally variable based on lake size. Nutrient concentrations in lakes are influenced by the surrounding landscape; and lakes that drain agricultural areas may be likely to have higher nutrient concentrations than those that drain forested areas. Nutrient concentrations can also be driven by lake depth and morphology. Shallower lakes tend to have higher nutrient concentrations because they are more biologically productive. Because many of the smaller program lakes likely mix and stratify several times throughout the summer (polymictic), they can exhibit higher nutrient concentrations than larger lakes that mix and stratify twice per year (dimictic). Although trophic classification is determined by specific nutrient concentrations, lake size is an important factor in determining the speed at which eutrophication can occur.

Summer profiles have indicated Beaver, Bootjack, Foy, Murray, Lake Blaine, and Jette were found to be anoxic at some depth. Eutrophic lakes that are covered in ice and snow with large quantities of decaying organic matter can exhibit substantial dissolved oxygen losses in the winter. When the organic matter supply for decomposition is large relative to the mass of dissolved oxygen available, there may be fish and macroinvertebrate kills (Kalff, 2002). In the winter of 2011/2012, Jette Lake had a winter fish kill due to dissolved oxygen depletion.

Lakes with hypoxic hypolimnia tend to be less than 10 m in depth, have high external phosphorus loading, and high algal biomass. Tally Lake is an exception and likely experiences hypoxia as a result of many complex processes including chemical oxidation of allochthonous humic matter in addition to the high wind energy required to mix this deep lake (Koopal, 2015 pers. comm.).

Program lakes are highly variable in terms of calcium concentration with a range from a low of 4 mg/L to 62 mg/L. Dreissinid mussels need calcium for shell development; however, calcium concentration is not the only relevant factor in determining whether a specific water body is suitable for mussel habitation. The calcium threshold is an indication for the potential of zebra or quagga mussel suitability among program lakes. Literature suggests the calcium threshold for Zebra mussel establishment can be <20 mg/L, however evidence suggests that long term survival, and establishment of zebra mussel colonies requires >28 mg/L, and 12-15 mg/L is the minimum concentration necessary for long term reproduction and growth (Cohen & Weinstein, 2008). This report classified waters with calcium < 20 mg/L as low risk for infestation, calcium 20 mg/L – 28 mg/L as moderate risk for infestation, and calcium >28 mg/L as high risk for zebra mussel infestation (Whittier *et al.*, 2008). Of the 53 monitoring sites that were sampled, 26 ranked as high risk, 20 ranked as moderate risk, and 7 ranked as low risk for calcium based risk assessment.

Alkalinity, pH, dissolved oxygen, and conductivity are other variables that determine suitability for long term survival of zebra/quagga mussels. Literature suggests that pH values below 7.3 are too acidic to support long term survival of zebra mussels, and levels between 7.4 and 9.4 are within the range for growth with 8.4 as ideal (Benson *et al*, 2017). Zebra mussels spawn between 12-18° C which is also the ideal temperature for larval development.

6.2 Study Limitations/Challenges

Through the years of the program, there have been challenges and unique situations that have occurred.

2011: The greatest challenge was the inability to begin the field season data collection on schedule because the Hydrolab MS5 that was expected to be used for the field season didn't arrive until October, and WLI's Hydrolab DS5 wasn't always available when needed. WLI's DS5 also needed to be shipped to Hach Environmental several times for repairs. However, Hydrolab profiles were taken on every lake at least once and were taken twice on 24 of the 42 lakes. The program now has the Hydrolab MS5 (and has back use of WLI's DS5).

2012: No summer Hydrolab profiles were taken on any of the Flathead Lake sites due to equipment maintenance issues.

2013: No fall Hydrolab profiles were collected on Flathead Lake, and water chemistry samples were not collected at three of the Flathead Lake sites. Fall Hydrolab profiles were collected at all eight Flathead Lake monitoring sites.

2014: Unseasonably cold weather caused many of the program lakes to freeze in early November. Because of this, fall Hydrolab profiles were not collected on some of the lakes.

2015: Hydrolab equipment repairs and fires prevented summer sampling on some of the program lakes.

2018: The transition from the NCSS software suite to using the open source computer language R has benefits in preparation of figures and databases and ensures quality control and assurance, but as with any conversion of database platforms, it created some graphic inconsistencies. In 2019, we will reconcile changes in graphic output and style.

General limitations: Water chemistry samples are only collected once each year at one site and Hydrolab profiles are only collected once each year at each lake limiting the spatial and temporal data. In addition, each lake is affected by seasonal variability. Data are intended to provide an inter-annual snapshot of summer conditions, hence the importance for consistency in sampling between mid-July and mid-August. The Carlson TSI also only includes one data point for each year except for Secchi depth, wherein all depths collected for the year are averaged then calculated in the TSI equation.

7.0 RECOMMENDATIONS

We recommend more intensive sampling on lakes with unique biological and chemical characteristics, those with native focal fish species, or lakes that have direct surface connection to those waters. Examples include Upper and Lower Stillwater, Upper Whitefish, Smith, Tally, Beaver, Swan, Lindbergh, Holland, Foy's Lake, and Flathead.

Given the increasing pressures on Montana's lakes from AIS, climate change, and increased human development, the need to gather consistent, accurate information on conditions from Montana's lakes has

never been greater. Lake volunteer water quality and AIS monitoring programs such as this one are important models for other regions and lake associations. With this in mind, WLI is initiating a program with funds from the Bureau of Reclamation and distributed through the Upper Columbia Conservation Commission (UC3) that will identify gaps in geographical coverage of lakes in the Upper Columbia river region and engage additional watershed, natural resource, and volunteer groups and citizens in long-term stewardship of basin lakes and reservoirs. This program will train Upper Columbia stakeholders in plankton tow net sampling techniques, visual identification of aquatic invasive plant species, and collecting water quality measurements. It will also provide those stakeholders with the necessary monitoring and decontamination equipment as well as an outreach messaging consistent with FWP's Clean, Drain, Dry campaign.

A number of other groups in northwest Montana (Swan Valley Connections, Blackfoot Challenge, Flathead Lakers, Flathead Lake Biological Station) support existing lake volunteer water quality and AIS monitoring programs. A large part of this project will be to enhance coordination and support among existing monitoring partners within the Upper Columbia Basin as well as supporting and outfitting new monitoring efforts. Incorporating additional partner groups adds to the volunteer base in the region and may allow more lakes to be monitored that are a high priority but are not currently sampled. WLI will create a new website highlighting all participating partner groups, a map of all lakes monitored and by whom, sampling and decontamination protocols, and background resources for volunteers.

This program relies on volunteers and project partners for long term success. Continued support is essential for the monitoring efforts to provide scientific data, education and outreach, and AIS early detection monitoring for the currently monitored forty-one lakes in northwest Montana. This information will help citizens and resource managers navigate the complex resource management decision making process.

Lastly, WLI is under contract with DEQ in 2019/2020 to upload data from the NWMTLVMN into DEQ's MT-eWQX EQUIS database. After completion of the upload, data from program lakes will be available to more users in order to better our overall understanding of lake resources.

8.0 LITERATURE CITED

- Benson, A.J., Raikow, D., Larson, J., Fusaro, A., & Bogdanoff, A.K. 2017. *Dreissena polymorpha*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL.
- Cohen, A.N., & Weinstein, A. 2001. Zebra Mussel's Calcium Threshold and Implications for its Potential Distribution in North America. San Francisco Estuary Institute, Richmond, CA. 44pp.
- Ellis, B.K. & J. A. Craft. 2008. Trophic Status and Trends in Water Quality for Volunteer Monitoring Program Lakes in Northwestern Montana, 1993-2007. FLBS Report #200-08. Prepared for Flathead Basin Commission, Kalispell, Montana by Flathead Lake Biological Station, The University of Montana, Polson, MT. 43pp.
- Kalff, J. 2002. *Limnology: Inland Water Ecosystems*. Prentice Hall. Upper Saddle River, New Jersey. 59pp.
- Koopal, M. 2014. Swan Lake Water Quality Investigation. Prepared for Swan Lakers, Kootenai Lodge Estates, and Swan Ecosystem Center. Whitefish Lake Institute, Whitefish, MT.
- Koopal, M. 2015. Personal Communication. WLI.
- NALMS management guide for lakes and reservoirs. North American Lake Management Society, P.O. Box 5443, Madison, WI, 53705-5443, USA (<http://www.nalms.org>). Accessed February 10th, 2012.
- Montana Fisheries Information System (MFISH). Montana Fish, Wildlife and Parks. 2009. <<http://fwp.mt.gov/fishing/mfish/default.aspx>>. Accessed January 10th, 2012.
- Reller, M. Used by permission of Bathymetry by Constellation Services; Copyright Protected 2004 – 2014; Not for Reproduction All Rights Reserved MarksLakeMaps.com.
- Wachsmuth, J. L. 2010. Developing a Citizen Volunteer Lake Monitoring Program for the Flathead Basin in Northwestern Montana for the early Detection and Prevention of the Introduction of Zebra Mussels and other Aquatic Invasive Species (AIS). University of Denver University College Capstone Project for Master of Applied Science.
- Wetzel, R.G. 2001. *Limnology*. (3rd edition). Academic Press, New York, NY. 1066 pp.
- Whitefish Lake Institute. 2011. Sampling and Analysis Plan. Northwest Montana Lakes Volunteer Monitoring Network (NWMTLVMN).
- Whittier, T. R., *et al.* *Frontiers in Ecology and the Environment, A Calcium-Based Invasion Risk Assessment for Zebra and Quagga Mussels*. *Front Ecol Environ* 2008; doi:10.1890/070073.

9.0 MAP SOURCES CITED

Abbot: *Google Maps*
Ashley: *MFWP*
Bailey: *Google Maps*
Beaver: *MFWP*
Big Therriault: *MFWP*
Blaine: *MFWP*
Blanchard: *MFWP*
Bootjack: *MFWP*
Dickey: *Mark Reller; Constellation Services*
Dollar: *MFWP*
Echo: *Mark Reller; Constellation Services*
Fish: *Google Maps*
Five: *MFWP*
Flathead: *Mark Reller; Constellation Services*
Foys: *MFWP*
Glen: *MFWP*
Halfmoon: *MFWP*
Hanson-Doyle: *Google Maps*
Holland: *Mark Reller; Constellation Services*
Jette: *Google Maps*
Lake of the Woods: *MFWP*
Lindbergh: *Mark Reller; Constellation Services*
Little Bitterroot: *Mark Reller; Constellation Services*
Loon: *MFWP*
Lost Coon: *Google Maps*
Lower Stillwater: *Mark Reller; Constellation Services*
Mary Ronan: *Mark Reller; Constellation Services*
McGilvray: *Google Maps*
Murphy: *Mark Reller; Constellation Services*
Murray: *MFWP*
Peterson: *Google Maps*
Rogers: *MFWP*
Skyles: *MFWP*
Sophie: *Mark Reller; Constellation Services*
Spencer: *MFWP*
Swan: *Mark Reller; Constellation Services*
Tally: *Mark Reller; Constellation Services*
Tetrault: *Google Maps*
Upper Stillwater: *Mark Reller; Constellation Services*
Upper Whitefish: *MFWP*
Whitefish: *Mark Reller; Constellation Services*

10.0 LIST OF ACRONYMS

AIS:	Aquatic Invasive Species
Chl (a):	Chlorophyll (a)
COC:	Chain of Custody
DEQ:	Department of Environmental Quality
DS5:	Data Sonde 5
eDNA:	Environmental Deoxyribonucleic Acid
EPA:	Environmental Protection Agency
EWM:	Eurasian Watermilfoil
FBC:	Flathead Basin Commission
FLBS:	Flathead Lake Biological Station
GPS:	Global Positioning System
HDPE:	High Density Polyethylene
LDO:	Luminescent Dissolved Oxygen
MDA:	Montana Department of Agriculture
MFWP:	Montana Fish, Wildlife and Parks
MS5:	Mini Sonde 5
NWMTLVMN:	Northwest Montana Lakes Volunteer Monitoring Network
ORP:	Oxidation Reduction Potential
pH:	Parts Hydrogen
SAP:	Sampling and Analysis Plan
SpC:	Specific Conductivity
TDS:	Total Dissolved Solids
TP:	Total Phosphorus
TPN:	Total Persulfate Nitrogen
TSI:	Trophic State Index
VLMP:	Volunteer Lakes Monitoring Program
WLI:	Whitefish Lake Institute

